

The high frequency antigen KANNO is located on prion protein, encoded by the *PRNP* gene, as a new blood group system

Kenichi Ogasawara



*Japanese Red Cross Society
Central Blood Institute, Blood Service Headquarters*

Characteristics of anti-KANNO

- **First example of anti-KANNO was reported in 1991**
- **First case of non-Japanese individual with anti-KANNO was reported in 2018 (Jones et al. ISBT Toronto)**



Characteristics of anti-KANNO

- Against unknown HFA (reactive with K_o , Jr(a-), Rh_{null} ...)
- Like a HTLA antibody
- Mainly detected in female with pregnancy history
- Clinical significance of anti-KANNO is unknown

Incompatible transfusion: 7cases

Pregnant women: 15 cases



No cases showed
HTR or HDFN

(Kawabata et al. *Transfus Med Rev* 2014; 28: 23-8)



Characteristics of KANNO antigen

- Sensitive to proteases Ficin, Trypsin, α -Chymotrypsin...
- Resistant to disulphide bond reducing agents AET and DTT
- KANNO- frequency is 0.44% (10 in 2,260)



What is the carrier molecule?

- IP, blotting, and MAIEA assays were failed...
- Genome-based approaches to identify the causal gene of KANNO antigen (Omae et al. *Transfusion* 2019)



Genome-Wide Human SNP array 6.0 (Affymetrix)

Genome-Wide Association Study (GWAS)


→ 4 KANNO- individuals vs. 415 healthy Japanese

Whole-Exome Sequencing (WES)

Sanger Sequencing




What is the carrier molecule?





Significant association on
Chromosome 20p13
rs6116471

c.655G>A (p.Glu219Lys)





PRNP genotype of the 4 probands and their family members

c.655G>A rs1800014



Monoclonal antibody-specific immobilization of erythrocyte antigen (MAIEA) assay





- KANNO antigen is on the prion protein
- Confirmed by transfection and expression study using CHO-K1



The *PRNP**655A frequencies in the ExAC database and in the Tohoku region

GG GA AA




Population	Allele Frequency (c.655G>A, rs1800014)	
South Asian	4.11%	(677 in 16,472)
East Asian	4.03%	(348 in 8,642)
Latino	0.19%	(22 in 11,560)
African	0.03%	(3 in 10,374)
European	0.004%	(3 in 66,660)
Tohoku (Japan)	5.80%	(58 in 1,000)



Correlation between KANNO phenotype and *PRNP* genotype

Phenotype	Agglutination strength	Number of samples*	c.655 genotype		
			GG	GA	AA
KANNO+	(2+-3+)	100	89	11	0
	(w+)	12	1	11	0
KANNO—	(0)	10	0	0	10

*Obtained from blood donors living in the Kanto-Koshinetsu region



Japanese Red Cross Society
Central Blood Institute, Blood Service Headquarters

Summary

- **Anti-KANNO may be stimulated by pregnancy or by transfusion**
- **Anti-KANNO appears to be clinically insignificant**
- **HFA KANNO is located on prion protein encoded by the *PRNP* gene**
- **Recessive inheritance of KANNO- is caused by the *PRNP**655A with c.655G>A (p.Glu219Lys) mutation**
- **The *PRNP**655A allele is more frequent in Asians than in other populations**



Thank you for your attention!



THE UNIVERSITY OF TOKYO

Yusuke Omae

Katsushi Tokunaga




FUKUSHIMA
MEDICAL
UNIVERSITY

Mayumi Takeuchi

Kinuyo Kawabata

Ikuo Wada

Hitoshi Ohto



日本赤十字社
Japanese Red Cross Society

Shoichi Ito

Kazumi Isa

Kenichi Ogasawara

Akira Oda

Sayaka Kaito

Hatsue Tsuneyama

Makoto Uchikawa